

Ukrainian Academy of Sciences  
Institute for Condensed Matter Physics

Preprint  
ICMP-94-9E

O.Derzhko, T.Krokhmalskii

Statistical properties of random spin- $\frac{1}{2}$   $XY$  chains

L'viv-1994

O.Derzhko, T.Krokhmalskii

Statistical properties of random spin- $\frac{1}{2}$   $XY$  chains

A numerical approach to the study of equilibrium statistical properties of spin- $\frac{1}{2}$   $XY$  chains is suggested. The approach is illustrated by the examining of influence of disorder on transverse dynamical susceptibility of spin- $\frac{1}{2}$  Ising chain in random transverse field.

©1994 Institute for Condensed Matter Physics

One-dimensional spin- $\frac{1}{2}$   $XY$  models are among the few many-body problems for which a lot of exact results can be obtained. This is due to the fact that the Hamiltonian of the system can be rewritten via Jordan-Wigner transformation in terms of non-interacting fermions [1]. In spite of this there are rather limited results for time-dependent correlations between longitudinal spin components. More difficult to deal with are the random versions of the model. Apart from academic importance such results are used in the description of a large number of quasi-one-dimensional ferroelectric and magnetic compounds such as  $Cs(H_{1-x}D_x)_2PO_4$ ,  $PbH_{1-x}D_xPO_4$ ,  $PrCl_3$  etc. In our previous paper [2] we suggested a new numerical approach for evaluation of thermodynamical properties and both equal-time and different-time spin correlation functions for spin- $\frac{1}{2}$  non-uniform anisotropic  $XY$  chain in transverse field with the Hamiltonian

$$H = \sum_{j=1}^N \Omega_j s_j^z + \sum_{j=1}^{N-1} (J_j^{xx} s_j^x s_{j+1}^x + J_j^{yy} s_j^y s_{j+1}^y). \quad (1)$$

This method allows to study the random versions of the model (1) assuming that  $\dots, \Omega_j, \dots; \dots, J_j^{xx}, \dots; \dots, J_j^{yy}, \dots$  vary randomly from site to site according to the given probability distribution. The purpose of the present paper is to describe in more detail the proposed approach and to apply it for examining the influence of disorder on transverse dynamical susceptibility of spin- $\frac{1}{2}$  Ising chain in random transverse field. Such results, up to the best authors' knowledge, are obtained for the first time.

In order to obtain the equilibrium statistical properties of the system in question (1) one should first perform Jordan-Wigner transformation from the raising and lowering operators  $s_j^\pm \equiv s_j^x \pm i s_j^y$  to Fermi operators  $c_1 = s_1^-, c_j = \prod_{n=1}^{j-1} (-2s_n^z) s_j^-, j = 2, \dots, N$ ,  $c_1^+ = s_1^+, c_j^+ = \prod_{n=1}^{j-1} (-2s_n^z) s_j^+, j = 2, \dots, N$  and

then to diagonalize the obtained quadratic form by a canonical transformation  $\eta_k = \sum_{j=1}^N (g_{kj}c_j + h_{kj}c_j^+)$  with the outcome  $H = \sum_{k=1}^N \Lambda_k(\eta_k^+ \eta_k - \frac{1}{2})$ . The elementary excitation spectrum  $\Lambda_k$  and the coefficients  $g_{kn} \equiv (\Phi_{kn} + \Psi_{kn})/2$ ,  $h_{kn} \equiv (\Phi_{kn} - \Psi_{kn})/2$  are determined from

$$\begin{aligned} \Lambda_k \Phi_{kn} &= \sum_{j=1}^N \Psi_{kj}(\mathbf{A} + \mathbf{B})_{jn}, \quad \Lambda_k \Psi_{kn} = \sum_{j=1}^N \Phi_{kj}(\mathbf{A} - \mathbf{B})_{jn}; \\ A_{ij} &\equiv \Omega_i \delta_{ij} + \frac{J_i^{xx} + J_i^{yy}}{4} \delta_{j,i+1} + \frac{J_{i-1}^{xx} + J_{i-1}^{yy}}{4} \delta_{j,i-1}, \\ B_{ij} &\equiv \frac{J_i^{xx} - J_i^{yy}}{4} \delta_{j,i+1} - \frac{J_{i-1}^{xx} - J_{i-1}^{yy}}{4} \delta_{j,i-1}. \end{aligned} \quad (2)$$

Eqs. (2) reduce to standard problems

$$\begin{aligned} \Lambda_k^2 \Phi_{kn} &= \sum_{j=1}^N \Phi_{kj}[(\mathbf{A} - \mathbf{B})(\mathbf{A} + \mathbf{B})]_{jn}, \\ \Lambda_k^2 \Psi_{kn} &= \sum_{j=1}^N \Psi_{kj}[(\mathbf{A} + \mathbf{B})(\mathbf{A} - \mathbf{B})]_{jn} \end{aligned} \quad (3)$$

for  $N \times N$  five diagonal banded matrices  $(\mathbf{A} \mp \mathbf{B})(\mathbf{A} \pm \mathbf{B})$ .

Next it is necessary to express the quantities of interest through  $\Lambda_k$ ,  $\Phi_{kj}$ ,  $\Psi_{kj}$ . One immediately notes that the evaluation of partition function  $Z(\beta, N) = \prod_{k=1}^N 2 \cosh \frac{\beta \Lambda_k}{2}$  involves all  $\Lambda_k^2$ . Thus, the knowledge of eigenvalues of matrices  $(\mathbf{A} \mp \mathbf{B})(\mathbf{A} \pm \mathbf{B})$  permits one to obtain the thermodynamical properties of the model in question (1). Somewhat more complicated is the investigation of spin correlation functions. Reformulating spin operators in terms of operators  $\varphi_j^\pm \equiv c_j^\pm \pm c_j$  (they are the known linear combinations of operators  $\eta_k, \eta_k^+$ , namely,  $\varphi_j^+ = \sum_{m=1}^N \Phi_{mj}(\eta_m^+ + \eta_m)$ ,  $\varphi_j^- = \sum_{m=1}^N \Psi_{mj}(\eta_m^+ - \eta_m)$ ), that is,  $s_j^x = \prod_{n=1}^{j-1} (\varphi_n^+ \varphi_n^-) \varphi_j^+ / 2$ ,  $s_j^y = \prod_{n=1}^{j-1} (\varphi_n^+ \varphi_n^-) \varphi_j^- / (2i)$ ,  $s_j^z = -\varphi_j^+ \varphi_j^- / 2$ , noting that the calculation of spin correlation function reduces to exploiting Wick-Bloch-de Dominicis theorem and calculating the elementary

contractions

$$\begin{aligned}
\langle \varphi_j^+(t) \varphi_m^+ \rangle &= \sum_{p=1}^N \Phi_{pj} \Phi_{pm} \frac{\cosh(i\Lambda_p t - \frac{\beta\Lambda_p}{2})}{\cosh \frac{\beta\Lambda_p}{2}}, \\
\langle \varphi_j^+(t) \varphi_m^- \rangle &= \sum_{p=1}^N \Phi_{pj} \Psi_{pm} \frac{\sinh(-i\Lambda_p t + \frac{\beta\Lambda_p}{2})}{\cosh \frac{\beta\Lambda_p}{2}}, \\
\langle \varphi_j^-(t) \varphi_m^+ \rangle &= - \sum_{p=1}^N \Psi_{pj} \Phi_{pm} \frac{\sinh(-i\Lambda_p t + \frac{\beta\Lambda_p}{2})}{\cosh \frac{\beta\Lambda_p}{2}}, \\
\langle \varphi_j^-(t) \varphi_m^- \rangle &= - \sum_{p=1}^N \Psi_{pj} \Psi_{pm} \frac{\cosh(i\Lambda_p t - \frac{\beta\Lambda_p}{2})}{\cosh \frac{\beta\Lambda_p}{2}}, \tag{4}
\end{aligned}$$

one finds that  $\Phi_{kj}$ ,  $\Psi_{km}$  for all  $\Lambda_k$  are involved in final expressions for both equal-time and different-time spin correlation functions. Therefore, the study of equilibrium statistical properties may be performed numerically in a following way: first, to solve the eigenvalues and eigenvectors problem for matrix  $(\mathbf{A} + \mathbf{B})(\mathbf{A} - \mathbf{B})$  (3) obtaining in result  $\Lambda_k^2$  and  $\Psi_{kj}$ , and second, having  $\Psi_{kj}$  and  $\Lambda_k = \sqrt{\Lambda_k^2}$ , to find  $\Phi_{kj}$  from Eq.(2); thermodynamics and spin correlation functions are expressed via the sought quantities. It should be stressed that here appears only the eigenvalues and eigenvectors problem for  $N \times N$  five diagonal banded matrix that is the remarkable peculiarity of the system in question (1). This is a key difference of our approach in comparison with other finite-chain calculations (reported e.g. in [3]) where the numerical diagonalization of the Hamiltonian (1) in the Hilbert space of dimension  $2^N$  is performed (see also [4, 5]).

We end up with presenting the results of numerical calculation of transverse dynamical susceptibility  $\chi_{zz}(\omega, \omega) \equiv \sum_n e^{i\omega n} \int_0^\infty dt e^{i(\omega + i\epsilon)t} \frac{1}{i} \langle [s_j^z(t), s_{j+n}^z] \rangle$  for spin- $\frac{1}{2}$  Ising chain ( $J_j^{xx} = J$ ,  $J_j^{yy} = 0$ ) in random transverse field  $\Omega_j$  with probability distribution density  $p(\dots, \Omega_j, \dots) = \prod_{j=1}^N [x\delta(\Omega_j) + (1-x)\delta(\Omega_j - \Omega)]$ . Acting as described above one finds that  $4 \langle s_j^z(t) s_{j+n}^z \rangle = \langle$

$$\varphi_j^+(t)\varphi_j^-(t)\varphi_{j+n}^+\varphi_{j+n}^- \Rightarrow \langle \varphi_j^+ \varphi_j^- \rangle \langle \varphi_{j+n}^+ \varphi_{j+n}^- \rangle - \langle \varphi_j^+(t)\varphi_{j+n}^+ \rangle \langle \varphi_j^-(t)\varphi_{j+n}^- \rangle + \langle \varphi_j^+(t)\varphi_{j+n}^- \rangle \langle \varphi_j^-(t)\varphi_{j+n}^+ \rangle, \langle s_{j+n}^z s_j^z(t) \rangle \Rightarrow \langle s_{j+n}^z(-t) s_j^z \rangle.$$

Computing then numerically the elementary contractions (4) involved in the spin correlation functions  $\langle s_{100}^z(t) s_{100+n}^z \rangle$ ,  $\langle s_{100+n}^z(-t) s_{100}^z \rangle$  for the random chain of 200 spins with  $J = -2$ ,  $\Omega = 1$  at  $\beta = 20$ , performing numerically the integration over  $t$  and summation over  $n$ , and averaging over the realization (typically few hundred) we have obtained the results some of which are presented in Fig.1. They agree with generally known results for perfect (non-random) cases of Ising model when  $x = 1$  and Ising model in transverse field when  $x = 0$  derived for the first time with periodic boundary conditions imposed in [6]. As the concentration of the sites with transverse field  $\Omega$  increases  $\chi_{zz}(\mathfrak{x}, \omega)$  rebuilds from the Ising-like type behaviour to the behaviour inherent to Ising model in transverse field. The divergence of  $\text{Re}\chi_{zz}(\mathfrak{x}, \omega)$  at  $\omega = 2$  becomes finite, decreases, loses its symmetry and develops into a smooth curve in the limiting case  $x = 0$ . The  $\delta$ -singularity at  $\omega = 2$  of  $\text{Im}\chi_{zz}(\mathfrak{x}, \omega)$  transforms into few sharp cusps, loses its symmetry moving slightly to the right and also develops into a smooth curve at  $x = 0$ . The study of the dependence of presented results on number of spins in the chain, on upper limit in integration over  $t$ , on value of  $\epsilon$ , on number of terms in sum over  $n$  and on number of random realizations were performed carefully. It should be noted that the obtained susceptibilities exhibits a lot of structure (especially at small  $x$ ) that requires further study.

Since a complete solution for some pair time-dependent correlation functions even in perfect spin- $\frac{1}{2}$  XY chains has never been found, the numerical calculations seem to be of much use in deriving new results here. Some rigorous results on longitudinal time-dependent correlation function were de-

rived numerically for perfect one-dimensional spin- $\frac{1}{2}$  transverse Ising model at critical field and isotropic  $XY$  model in zero field at  $\beta = \infty$  in [7], however, within quite different approach. The suggested method, that gives exact results for finite spin- $\frac{1}{2}$   $XY$  chains, appears to be of great use in study of dynamical properties and random versions of systems considered. The derived conclusions may be useful for interpretation of observable data in dynamical experiments on quasi-one-dimensional compounds that can be described in frames of model (1).

The present paper was presented as a poster at International Conference on Magnetism 1994 (22-26 August, 1994, Warsaw, Poland) (EP.227) [8]. The authors would like to express their gratitude to Prof. L.L.Gonçalves and other participants of ICM'94 for interest in this paper and discussions.

## References

- [1] E.Lieb, T.Schultz, D.Mattis, *Ann.Phys.* **16**, 407 (1961).
- [2] O.V.Derzhko, T.Ye.Krokhmal'skii, *Visn.L'viv.Univ., ser.fiz.* 26, 47 (1993) (in Ukrainian).
- [3] M.D'Iorio, U.Glaus, E.Stoll, *Solid State Commun.* **47**, 313 (1983).
- [4] P.M.Richards, F.Carboni, *Phys.Rev.B* **5**, 2014 (1972).
- [5] J.P.Groen, T.O.Klaassen, N.J.Poulis, G.Müller, H.Thomas, H.Beck, *Phys.Rev.B* **22**, 5369 (1980).
- [6] Th.Niemeijer, *Physica* **36**, 377 (1967).
- [7] G.Müller, R.E.Shrock, *Phys.Rev.B* **29**, 288 (1984).

- [8] O.Derzhko, T.Krokhmalskii, *International Conference on Magnetism 1994, 22-26 August, 1994, Warsaw, Poland. Programme and Abstracts*, 848 (1994).



Fig. 1. Frequency-dependent real and imaginary parts of transverse susceptibility  $\chi_{zz}(\omega)$  for Ising chain in random transverse field at  $\omega = 0$  for different values of concentration  $x$ ;  $\epsilon = 0.01$ .

Fig. 1. Frequency-dependent real and imaginary parts of transverse susceptibility  $\chi_{zz}(\omega)$  for Ising chain in random transverse field at  $\omega = 0$  for different values of concentration  $x$ ;  $\epsilon = 0.01$ .

Address:

Institute for Condensed Matter Physics

1 Svientsitskii St., L'viv-11, 290011, Ukraine

Tel.: (0322)761054

Fax: (0322)761978

E-mail: [icmp@sigma.icmp.lviv.ua](mailto:icmp@sigma.icmp.lviv.ua)